

On the origin of the Schechter-like mass function of young star clusters in disk galaxies

P. Lieberz¹, P. Kroupa^{1,2}

¹ *Helmholtz Institut für Strahlen- und Kernphysik (HISKP), Universität Bonn, Nussallee 14–16, D-53121 Bonn, Germany*

² *Charles University in Prague, Faculty of Mathematics and Physics, Astronomical Institute, V Holešovičkách 2, CZ-180 00 Praha 8, Czech Republic*

Released 2002 Xxxxx XX

ABSTRACT

The mass function of freshly formed star clusters is empirically often described as a power law. However the cluster mass function of populations of young clusters over the scale of a galaxy has been found to be described by a Schechter-function. Here we address this apparent discrepancy. We assume that in an annulus of an isolated self-regulated radially-exponential axially-symmetric disk galaxy, the local mass function of very young (embedded) clusters is a power law with an upper mass limit which depends on the local star formation rate density. Radial integration of this mass function yields a galaxy-wide embedded cluster mass function. This integrated embedded cluster mass function has a Schechter-type form, which results from the addition of many low mass clusters forming at all galactocentric distances and rarer massive clusters only forming close to the center of the galaxy.

Key words: stars: formation – stars: luminosity function, mass function – stars: statistics – galaxies: star clusters – galaxies: stellar content

1 INTRODUCTION

The freshly-formed stellar mass of a galaxy resides in its embedded star clusters (Lada & Lada 2003; Kroupa 2005; Megeath et al. 2016; Meingast et al. 2016), which can synonymously be referred to as (about 1 pc extended, 1 Myr duration) space-time correlated star-formation events. i.e. essentially in molecular cloud clumps. The investigation of embedded clusters helps us to understand the build up of the stellar populations in galaxies. Embedded clusters are still fully or partially enshrouded in gas and dust, and represent the earliest stage in the life-time of a formed star cluster. They may be the forerunners to the open clusters and are therefore valuable probes for cluster formation (Lada & Lada 2003).

Observations indicate that the masses of the embedded star clusters in galaxies follow a particular distribution function, the embedded cluster mass function (ECMF). Locally it is typically found to be a power-law, while galaxy-wide observation reveal a Schechter-like turn-down (Gieles et al. 2006). This has also been observed for old, massive globular clusters (see e.g. Jordan et al. 2007; Burkert & Smith 2000; Parmentier & Gilmore 2007). Massive clusters experience limited secular mass loss and are therefore still a reasonable indicator for the initial cluster mass function (Baumgardt & Makino 2003).

The focus of this paper is on embedded clusters in the disks of isolated late-type galaxies. Ideally the disk can be seen as a purely self-regulated axis-symmetric system, such that it is possible to investigate the ECMF only radially over the area of the disk galaxy to explain the difference between the local and integrated distribution¹.

In this paper we present for the first time analytical approaches to the galaxy wide ECMF and also rewrite the dependency on the galactocentric distance into a mass dependency so that we arrive at a new relation between the number and the mass of star clusters. Hereby we follow the ansatz that the galaxy-wide ECMF is the sum of all local ECMFs.

We constrain these models with observational data and show that the models account for the data quite well.

¹ The assumption of the galaxy being axis-symmetric is made for computational ease. The model developed here is equally valid in any galaxy in which the most massive cluster forming in a population of embedded clusters depends on the local gas density.

2 THE LOCAL EMBEDDED CLUSTER MASS FUNCTION

The local ECMF (LECMF or ξ_{lecl}) describes the surface number density of star clusters in the stellar mass interval $[M_{\text{ecl}}, M_{\text{ecl}} + dM_{\text{ecl}}]$ in an infinitesimally small surface area (dA) at a distance r from the center of the star-forming disk galaxy

$$\xi_{\text{lecl}}(M_{\text{ecl}}; r) dM_{\text{ecl}} dA = dN_{\text{ecl}}, \quad (1)$$

where dN_{ecl} is the number of embedded clusters in dM_{ecl} and dA .

In general the LECMF has the form of a power law, as derived from observations (Lada & Lada 2003)

$$\xi_{\text{lecl}}(M_{\text{ecl}}; r) = K(r) M_{\text{ecl}}^{-\beta}, \quad (2)$$

where M_{ecl} is the stellar mass of the embedded cluster at “birth”² and $K(r)$ the normalization constant and β the power law index. By observation β was found to be between $\beta = 1.5$ and $\beta = 2.5$ (Weidner, Kroupa & Larsen 2004, hereafter WKL, and references therein). $K(r)$ has to be estimated using normalization conditions.

The normalization condition is that the entire freshly formed stellar mass of a galactic region has to be in embedded clusters. So integrating the mass over all clusters (from the lower mass limit $M_{\text{ecl},\text{min}}$ to the locally upper mass limit $M_{\text{U},\text{loc}}(r)$) obtains the freshly formed stellar mass. In the case of an infinitesimally small surface area dA this obtains the embedded cluster mass surface density. This embedded cluster mass surface density is defined as the star formation rate density $\Sigma_{\text{SFR}}(r)$ multiplied with the time-scale, δt , which is the time over which a population of embedded clusters forms. $\Sigma_{\text{SFR}}(r)$ is defined as

$$\Sigma_{\text{SFR}}(r) = \frac{d\text{SFR}}{dA}, \quad (3)$$

with SFR being the star formation rate in the whole galaxy. The total embedded cluster mass density is obtained by multiplying $\Sigma_{\text{SFR}}(r)$ with δt (Schulz, Pflamm-Altenburg & Kroupa 2015; WKL):

$$\Sigma_{\text{SFR}}(r) \delta t = \int_{M_{\text{ecl},\text{min}}}^{M_{\text{U},\text{loc}}(r)} M'_{\text{ecl}} \xi_{\text{lecl}}(M'_{\text{ecl}}; r) dM'_{\text{ecl}}. \quad (4)$$

$M_{\text{ecl},\text{min}}$ is assumed to be about $5 M_{\odot}$, corresponding to the smallest groups of embedded stars observed (Kroupa & Bouvier 2003, Kirk & Myers 2012), and is here assumed to be a constant. δt is roughly 10 Myr, as deduced by Egusa, Sofue & Nakanishi (2004). This is the time it takes for the interstellar medium to transform into a new population of stars and is essentially the lifetime of molecular clouds (Fukui et al. 1999; Yamaguchi et al. 2001; Tamburro et al. 2008). It corresponds to the lifetime over which the embedded cluster mass function is fully populated (see the discussion in Kroupa et al. 2013 and Schulz et al. 2015).

It is noteworthy that to calculate the LECMF Eq. (4) is sufficient once $M_{\text{U},\text{loc}}(r)$ is known. The values for $K(r)$ are uniquely defined for different β . For $\beta \neq 2$,




Figure 1. An exemplary LECMF. Shown here is a LECMF from the distribution function given by Eq. (2) for a galactic region with $10^8 M_{\odot}$ of stellar mass. Further we assumed a $M_{\text{ecl},\text{min}}$ of $5 M_{\odot}$ and a $M_{\text{U},\text{loc}}(r)$ of $10^8 M_{\odot}$. The solid line shows the LECMF calculated analytically, whereas the points are the binned results of stochastic sampling.

$$K(r) = \frac{\Sigma_{\text{SFR}}(r) \delta t (2 - \beta)}{M_{\text{U},\text{loc}}^{2-\beta}(r) - M_{\text{ecl},\text{min}}^{2-\beta}}. \quad (5)$$

For the special case of $\beta = 2$,

$$K(r) = \frac{\Sigma_{\text{SFR}}(r) \delta t}{\ln(M_{\text{U},\text{loc}}(r)/M_{\text{ecl},\text{min}})}. \quad (6)$$

A visual verification that a LECMF calculated by this method is in agreement with a LECMF that was obtained by randomly drawing star clusters according to Eq. (2) is shown in Fig. 1.

The only uncertainty left is the exact form of $M_{\text{U},\text{loc}}(r)$. At the moment we can only state that the local upper mass limit has to be smaller than the cluster upper mass limit of the whole galaxy M_{U} : $M_{\text{U},\text{loc}}(r) \leq M_{\text{U}}$. Therefore additional research into this subject is necessary (see Sec. 3.1).

The LECMF with an observable non-infinitesimal surface area ΔA is discussed in Appendix A.

3 THE GALAXY-WIDE EMBEDDED CLUSTER MASS FUNCTION

In this section the scope of our analysis is to describe the form of the ECMF after integrating over the axis-symmetric galactic disk surface area. The previous formulae allow the normalisation of the LECMF at a fixed distance to the center of a galaxy.

The galaxy wide or integrated ECMF (IECMF or ξ_{iecl} , not to be confused with the LECMF ξ_{lecl} above) is defined as the number of star clusters in the stellar mass interval M_{ecl} to $M_{\text{ecl}} + dM_{\text{ecl}}$

$$\xi_{\text{iecl}}(M_{\text{ecl}}) dM_{\text{ecl}} = dN_{\text{ecl}}, \quad (7)$$

or

$$\begin{aligned} \xi_{\text{iecl}}(M_{\text{ecl}}) &= \int_0^{2\pi} \int_0^{R'(M_{\text{ecl}})} \xi_{\text{lecl}}(M_{\text{ecl}}; r) r dr d\phi \\ &= \int_0^{2\pi} \int_0^{R'(M_{\text{ecl}})} K(r) M_{\text{ecl}}^{-\beta} r dr d\phi. \end{aligned} \quad (8)$$

² “birth” describes here the idealised state when the entire population is existing directly prior to the expulsion of the gas.

with $R'(M_{\text{ecl}})$ being the maximal galactocentric distance at which an embedded cluster of mass M_{ecl} can form, assuming the maximum cluster mass decreases monotonically with increasing radial distance.

Is the IECMF still similar to a power-law function of the cluster mass? And if yes, does the power-law parameter β differ from the local one? These questions are addressed in the following.

The galaxy wide star formation rate (SFR)

$$\text{SFR} = \int_0^{2\pi} \int_0^{R_{\text{gal}}} \Sigma_{\text{SFR}} r \, dr d\phi, \quad (9)$$

with R_{gal} being the radius of the star forming area of the galaxy. In order to obtain ξ_{iecl} one can insert Eq. (4) into Eq. (9). The resulting equation (see also Kroupa et al. 2013) is:

$$\begin{aligned} \text{SFR} \, \delta t &= \int_0^{2\pi} \int_0^{R_{\text{gal}}} \int_{M_{\text{ecl},\min}}^{M_{\text{U},\text{loc}}(r)} M_{\text{ecl}} \xi_{\text{iecl}}(M_{\text{ecl}}; r) dM_{\text{ecl}} r \, dr d\phi \\ &= \int_0^{2\pi} \int_0^{R_{\text{gal}}} \int_{M_{\text{ecl},\min}}^{M_{\text{U},\text{loc}}(r)} K(r) M_{\text{ecl}}^{1-\beta} r \, dM_{\text{ecl}} dr d\phi. \end{aligned}$$

Eq. (10) is first an integration over the mass of the clusters and then over the area. It is possible to exchange the mass integral with the integral over the galactocentric distance by noting that in a given annulus the theoretical most massive cluster $M_{\text{U},\text{loc}}(r)$ depends on r , that is, we can invert this function to obtain the galactocentric distance, $R'(M_{\text{ecl}})$, at which the annulus contains a particular theoretical most massive cluster. In other words to exchange the positions of the r - and M_{ecl} -integration. Assuming that the mass of the theoretical most massive cluster possible, $M_{\text{U},\text{loc}}(r)$, decreases with the galactocentric distance r then the integration needs only to extend over the distances 0 to $R'(M_{\text{ecl}}) \leq R_{\text{gal}}$. Therewith at the low mass end the IECMF is an integral over 0 to R_{gal} , while at the high mass end large radii do not contribute:

$$\text{SFR} \, \delta t = \int_{M_{\text{ecl},\min}}^{M_{\text{U}}} \int_0^{2\pi} \int_0^{R'(M_{\text{ecl}})} K(r) M_{\text{ecl}}^{1-\beta} r \, dr d\phi dM_{\text{ecl}}. \quad (11)$$

Neither $K(r)$ nor $M_{\text{U},\text{loc}}(r)$ are known for a specific r . $K(r)$ depends on Σ_{SFR} , see Eq. (4). A direct relation between Σ_{SFR} and the position within the galaxy, assumed to be valid for a galaxy in self-regulated equilibrium, is taken from Pflamm-Altenburg & Kroupa (2008):

$$\begin{aligned} \Sigma_{\text{SFR}}(r) &= \frac{\text{SFR} \, e^{R_{\text{gal}}/r_d}}{2\pi r_d^2 \left(e^{R_{\text{gal}}/r_d} - \frac{R_{\text{gal}}}{r_d} - 1 \right)} e^{-r/r_d} \\ &\approx \frac{\text{SFR}}{2\pi r_d^2} e^{-r/r_d}. \end{aligned} \quad (12)$$

In this case r_d is the disc scale length. The equation is normalised in such a way that an integral over the whole area results in the total SFR. Therefore both sides of Eq. (12) can be multiplied with δt so that it is equal to Eq. (4) (in the following we will write $\text{SFR} \, \delta t$ as M_{tot} , the total stellar mass formed galaxy wide in time δt):

$$\int_{M_{\text{ecl},\min}}^{M_{\text{U},\text{loc}}(r)} M'_{\text{ecl}} \xi_{\text{iecl}}(M'_{\text{ecl}}) dM'_{\text{ecl}} = \frac{M_{\text{tot}}}{2\pi r_d^2} e^{-r/r_d}. \quad (13)$$

This gives a relation between $K(r)$ and $M_{\text{U},\text{loc}}(r)$. For $\beta \neq 2$:

$$K(r) = \frac{(2-\beta) M_{\text{tot}}}{2\pi r_d^2 (M_{\text{U},\text{loc}}^{2-\beta}(r) - M_{\text{ecl},\min}^{2-\beta})} e^{-r/r_d}. \quad (14)$$

For $\beta = 2$:

$$K(r) = \frac{M_{\text{tot}}}{2\pi r_d^2 \ln(M_{\text{U},\text{loc}}(r)/M_{\text{ecl},\min})} e^{-r/r_d}. \quad (15)$$

To get an unambiguous expression for $K(r)$ and $M_{\text{U},\text{loc}}(r)$ more constraints are needed. For this purpose we use a model (henceforth called the exponential model) based on an ansatz from Pflamm-Altenburg & Kroupa (2008). Other possible models, found to be not working as well as this one, are discussed in Appendix C.

3.1 Exponential Model

Pflamm-Altenburg & Kroupa (2008) propose the ansatz that the radial dependence of $M_{\text{ecl},\text{max},\text{loc}}(r)$ should have the same form as the radial dependence of the gas surface density $\Sigma_{\text{gas}}(r)$:

$$\Sigma_{\text{gas}}(r) = \Sigma_{\text{gas},0} e^{-\frac{r}{r_d}}, \quad (16)$$

with $\Sigma_{\text{gas},0}$ being the gas surface density at the center of the galactic disk. Thus

$$M_{\text{ecl},\text{max},\text{loc}}(r) = M_{\text{ecl},\text{max}} e^{-\frac{r}{r_d}}. \quad (17)$$

With this ansatz they were able to show that the $\text{H}\alpha$ radial cut-off in disk galaxies is naturally explained, given that star formation extends well beyond this cut-off radius.

As our model uses $M_{\text{U},\text{loc}}$ instead of $M_{\text{ecl},\text{max}}$, we modify Eq. (17) to

$$M_{\text{U},\text{loc}}(r) = M_{\text{U}} e^{-\frac{r}{r_d}}. \quad (18)$$

Inserting Eq. (18) into Eq. (13) with $\beta \neq 2$ results in a definite $K(r)$:

$$K(r) = \frac{(2-\beta) M_{\text{tot}}}{2\pi r_d^2 \left[\left(M_{\text{U}} e^{-\frac{r}{r_d}} \right)^{2-\beta} - M_{\text{ecl},\min}^{2-\beta} \right]} e^{-r/r_d}, \quad (19)$$

and for $\beta = 2$ in

$$K(r) = \frac{M_{\text{tot}}}{2\pi r_d^2 \ln \left[\left(M_{\text{U}} e^{-\frac{r}{r_d}} \right) / M_{\text{ecl},\min} \right]} e^{-r/r_d}. \quad (20)$$

For each r in the axis-symmetric disk galaxy there is a theoretical maximal cluster mass $M_{\text{U},\text{loc}}(r)$. Regarding the entire galaxy, each $M_{\text{U},\text{loc}}(r)$ is a theoretical possible cluster mass M_{ecl} . And in this relation r is the maximal galactocentric distance $R'(M_{\text{ecl}})$, at which a cluster of mass M_{ecl} can still be found. This is true for every M_{ecl} , $R'(M_{\text{ecl}})$ being the reverse function of $M_{\text{U},\text{loc}}(r)$. $R'(M_{\text{ecl}})$ is needed for Eq. (11),

$$R'(M_{\text{ecl}}) = -r_d \ln \left(\frac{M_{\text{ecl}}}{M_{\text{U}}} \right). \quad (21)$$

Now $\xi_{\text{iecl}}(M_{\text{ecl}})$ can be calculated:

$$\xi_{\text{iecl}}(M_{\text{ecl}}) = \int_0^{2\pi} \int_0^{R'(M_{\text{ecl}})} K(r) M_{\text{ecl}}^{-\beta} r \, dr d\phi. \quad (22)$$

This integration can only be solved numerically (because of the r -dependence in $K(r)$). The remaining free parameters (β and M_{U}) can be fixed using empirical data. WKL derived a fitting function for the mass of the most massive very young cluster in a galaxy ($M_{\text{vyc},\text{max}}$) depending on the SFR of the host galaxy:

$$M_{\text{vyc,max}} = k_{\text{ML}} \cdot \text{SFR}^{0.75(\pm 0.03)} \cdot 10^{6.77(\pm 0.02)}, \quad (23)$$

with SFR in M_{\odot}/yr and k_{ML} being the mass-to-light ratio in the photometric V-band, which can be assumed to be $0.0144 M_{\odot}/L_{V,\odot}$ for young (< 6 Myr) clusters. $M_{\text{vyc,max}}$ is a good approximation for the stellar mass of the most massive embedded star cluster of a galaxy, $M_{\text{ecl,max}}$.

To determine $M_{\text{ecl,max}}$ we use two conditions: first there is only one most massive cluster. Second the mass of the most massive cluster is $M_{\text{ecl,max}}$.

To implement the first condition we choose a mass interval between the upper mass limit M_U and $M_{\text{ecl,t}}$, with $M_{\text{ecl,t}}$ chosen in such a way that there is only one cluster between these limits:

$$1 = \int_{M_{\text{ecl,t}}}^{M_U} \xi_{\text{iecl}}(M'_{\text{ecl}}) dM'_{\text{ecl}}. \quad (24)$$

The second condition implies that the mass between these limits is $M_{\text{ecl,max}}$:

$$M_{\text{ecl,max}} = \int_{M_{\text{ecl,t}}}^{M_U} M'_{\text{ecl}} \xi_{\text{iecl}}(M'_{\text{ecl}}) dM'_{\text{ecl}}. \quad (25)$$

As $M_{\text{ecl,max}}$ depends on SFR it therewith follows that also M_U depends on SFR. Note that these conditions differ from those employed on previous occasions, as explained in Appendix B.

Using these we can find for every β a SFR- $M_{\text{ecl,max}}$ -curve that aligns with Eq. (23). In order to narrow β down, a fitting M_U is needed. M_U is supposed to be larger than any observed cluster mass but not so large that there would be unrealistic gaps between the mass of the most massive clusters and M_U . In the following we assume $r_d = 2.15$ kpc, as this is roughly the determined disk scale length of the Milky Way (Bovy & Rix 2013; Porcel et al. 1998). With that we find for $\beta = 2.3 \pm 0.1$ and $\delta t = 10$ Myr a SFR- $M_{\text{ecl,max}}$ -curve that aligns with Eq. (23) and resulting M_U that fulfils the above criteria (see also Fig. 2). This way we can give for every SFR a corresponding M_U . E.g. for $\text{SFR} = 1 M_{\odot}/\text{yr}$ we obtain $M_U = 400000 M_{\odot}$.

4 COMPARISON TO EMPIRICAL DATA

We already used empirical data in Sec. 3.1 to align the SFR- $M_{\text{ecl,max}}$ -curve of the exponential model with the empirical fit by WKL (Eq. 23) and in doing so constrained the free parameters (β and M_U) of the model. In the following we want to determine whether the model, using said constraints, is also in reasonable agreement with other observations.

Empirical data has been indicating that the galaxy-wide ECMF should be a Schechter-function (Gieles et al. 2006), i.e.

$$\xi_{\text{iecl,Schechter}}(M_{\text{ecl}}) = K' e^{-M_{\text{ecl}}/M_c} M_{\text{ecl}}^{-\beta'}. \quad (26)$$

³ In order to calculate $M_{\text{vyc,max}}$ from the observed absolute magnitude (M_V) the following formula has been used (Abdullah private communication):

$$M_V = 4.79 - 2.5 \log_{10} \frac{M_{\text{ecl,max}}}{k_{\text{ML}}},$$

with k_{ML} being the mass to light ratio.




Figure 2. The observational data was taken from WKL for $M_{\text{vyc,max}} < 2 \cdot 10^5 M_{\odot}$ and Randriamanakoto et al. (2013)³ for $M_{\text{vyc,max}} > 2 \cdot 10^5 M_{\odot}$ and shows the observed galaxy-wide most massive very young clusters $M_{\text{vyc,max}}$ (which are a good approximation for $M_{\text{ecl,max}}$) in dependence of SFR. Also plotted here is $M_{\text{ecl,max}}$ and M_U , as calculated using the model from Sec. 3.1 for $\beta = 2.3$ against the galaxy-wide SFR. All observed clusters should be below the M_U -line, which is mostly the case given the uncertainties.




Figure 3. A comparison between the IECMF acquired from the exponential model (the solid line) and the IECMF from the Schechter form (Eq. 26, the dashed line) for $\text{SFR} = 1 M_{\odot}/\text{yr}$ and a $\delta t = 10$ Myr. For the exponential model $\beta = 2.3$, while for the $\xi_{\text{iecl,Schechter}}$ model $\beta = 2.24$.

In this case M_c is the turn-down mass, at which the ECMF turns down. Our formalism results in a similar form for the ECMF as the Schechter form. This is shown in Fig. 3.

Note that the $\xi_{\text{iecl,Schechter}}$ model uses a different β than the exponential model. The exponential model has a sharper turn down than the Schechter formalism for the same β . Therefore β needs to be modified for a good fit. The smaller the SFR the bigger the divergence between the needed β form the exponential model and the Schechter form.

Up until now no theoretical formulation existed which allowed M_c to be predicted from properties of the galaxy.




Figure 4. Combined binned young cluster mass function for two galaxies (NGC 5236 and NGC 6946, data taken from Larsen 1999, Larsen 2009 and Larsen private communication). Also shown is the exponential model. For the single galaxies see Figs. 5 and 6.

With our formulation we can derive the turn-down mass in dependence of the exponential density profile of the galaxy.

Fitting the Schechter-function to the exponential model allows us to determine a relation between M_c and the SFR:

$$M_c = (85000 \pm 5000) \text{ SFR}^{(0.73 \pm 0.02)}. \quad (27)$$

Comparing this to Eq. (23) shows that these two equations are the same within the uncertainties. Therefore $M_c = M_{\text{ecl,max}}$ is at least a good approximation.

Next we compare the exponential model to actual empirical data. For this we confront the model with galaxy-wide observations of NGC 5236 and NGC 6946 (data taken from Larsen 1999, Larsen 2009 and Larsen private communication). The empirical data consists of a list of cluster masses, which were binned. To compare these data to the models we need the SFRs of the galaxies. Dopita et al. (2010) calculated a SFR of $2.76 M_\odot/\text{yr}$ for NGC 5236, whereas Hong et al. (2013) determined SFRs of 1.52 and $0.18 M_\odot/\text{yr}$, depending on the method used. For NGC 6946 Heesen et al. (2014) measured, depending on the method, a SFR of 4.6 ± 0.2 and $3.5 \pm 0.2 M_\odot/\text{yr}$. For our models we are therefore adopting $\text{SFR} = 1.5 M_\odot/\text{yr}$ for NGC 5236 and $\text{SFR} = 4 M_\odot/\text{yr}$ for NGC 6946. From Larsen (2009) we get the ages of the brightest and 5th brightest cluster. From these we can assume that $\delta t = 10 \text{ Myr}$ is a good approximation for the age. As $\xi_{\text{ecl}} \propto K \propto \delta t$ a change in δt does not change the overall behaviour of the function. A comparison of the data to the combined ξ_{ecl} calculated using the exponential model is shown in Fig. 4. The only difference between the IECMF models that has been applied here are the differing galaxy-wide SFRs. All other parameters are identical to the fit from Fig. 2. We can also compare the model to the individual galaxies (Figs. 5 and 6) but this has the disadvantage of having higher uncertainties. Nonetheless the model fits quite well, given the uncertainties.

The r -dependency of $M_{\text{U,loc}}(r)$ also allows a test of the exponential model. We compare the theoretical $M_{\text{U,loc}}-r$ dependence with the observed very young star clusters in M33 (data taken from




Figure 5. Binned young cluster mass function for NGC 5236 (data taken from Larsen 1999, Larsen 2009 and Larsen private communication). Also shown is the exponential model for $\text{SFR} = 1.5 M_\odot/\text{yr}$ and for $\text{SFR} = 2.76 M_\odot/\text{yr}$. See text for further details.




Figure 6. Binned young cluster mass function for NGC 6946 (data taken from Larsen 1999, Larsen 2009 and Larsen private communication). Also shown is the exponential model for $\text{SFR} = 4 M_\odot/\text{yr}$. See text for further details.

Pflamm-Altenburg, González-Lópezlira & Kroupa 2013) in Fig. 7. Fig. 7 shows the r -dependence of $M_{\text{U,loc}}$ for the model compared to observed very young star clusters. As $M_{\text{U,loc}}$ is the upper mass limit for star clusters at a given galactocentric distance we would expect that no observed cluster is heavier than it. But several observed ones are. Also using the observed SFR ($0.16 M_\odot/\text{yr}$, Skibba et al. 2011) in the empirical fit by Weidner (Eq. 23) results in a $M_{\text{vyc,max}}$ much smaller than several observed clusters. But taking into account that individual cluster masses have large uncertainties it becomes apparent that the exponential model fits quite well to the observed radially decreasing upper masses.




Figure 7. Comparison of the dependence on the galactocentric distance of $M_{U,loc}$ for the exponential model with observed very young clusters in M33 (Pflamm-Altenburg et al. 2013). The model uses $SFR = 0.16 M_{\odot}/yr$, $\delta t = 10$ Myr and $\beta = 2.31$. These values have been chosen so that the model fits the experimental data from the SFR - $M_{ecl,max}$ relation (Fig. 2).

5 CONCLUSION

In this paper we calculated for the first time the galaxy-wide integrated embedded cluster mass function for galaxies, which we assumed for computational ease to be axis-symmetric exponential disks, and showed that it has the form of a Schechter-like function, although locally the ECMF is a pure power-law.

To do that we first described an analytical solution of the LECMF. It was assumed there that the theoretical local upper cluster mass limit, $M_{U,loc}(r)$, depends on the galactocentric distance r .

Integrating the LECMF over the area of the star forming disk yields the galaxy-wide or integrated embedded cluster mass function (IECMF or ξ_{iecl}). For this purpose a model describing the r -dependence of $M_{U,loc}(r)$ was needed. The exponential model, introduced by Pflamm-Altenburg & Kroupa (2008), is found to be in agreement with observational data.

Even though the LECMF is a power law, the IECMF resembles a Schechter-like function, the reason being that the upper mass limit of the local power law is defined by $M_{U,loc}(r)$, which decreases with an increasing galactocentric distance.

Additionally given a locally estimated power-law index β for an ensemble of embedded clusters in a region in a galaxy, we have shown here that other regions elsewhere are expected to have different logarithmic ECMF, depending on the size and position of the region in the galaxy, and that the size of the region implies a Schechter-type turn-down of the embedded cluster mass function. The galaxy-wide ECMF, the integrated ECMF, IECMF, thus becomes a Schechter-type form.

All of this opens further possibilities for research. We assumed radially axis-symmetric disk galaxies but it is worth investigating whether a barred galaxy would cause significant changes. Also as has been stated, the exponential model depends on the theoretical upper limit for the mass of clus-

ters in a galaxy of a specific stellar mass. At the moment we can only give a lower limit to these values, which could underestimate them. Further research in the context of star and cluster formation should be able to give improved insights on the theoretical upper mass limits. Another step, which has to be done, would be to combine the stellar initial mass function (IMF) and the ECMF into the integrated galactic initial mass function (IGIMF) (Weidner et al. 2013, Recchi & Kroupa 2015), which is the galaxy-wide stellar initial mass function.

ACKNOWLEDGEMENTS

We thank the referee for helpful comments and Søren S. Larsen for providing us with valuable data regarding the galaxies NGC 5236 and NGC 6946. We are also grateful to Jan Pflamm-Altenburg for providing data on M33 and for useful discussions.

REFERENCES

- Baumgardt H., Makino J., 2003, MNRAS, 340, 227
- Bovy J., Rix H.-W., 2013, ApJ, 779, 115
- Burkert A., Smith G. H., 2000, ApJ, 542, L95
- Dopita M. A. et al., 2010, ApJ, 710, 964
- Egusa F., Sofue Y., Nakanishi H., 2004, PASJ, 56, L45
- Fukui Y. et al., 1999, PASJ, 51, 745
- Gieles M., Larsen S. S., Bastian N., Stein I. T., 2006a, A&A, 450, 129
- Heesen V., Brinks E., Leroy A. K., Heald G., Braun R., Bigiel F., Beck R., 2014, AJ, 147, 103
- Hong S., Calzetti D., Gallagher III J. S., Martin C. L., Conselice C. J., Pellerin A., 2013, ApJ, 777, 63
- Jordan A. et al., 2007, ApJS, 171, 101
- Kirk H., Myers P. C., 2012, ApJ, 745, 131
- Kroupa P., 2005, ESA Special Publ., 576, 629
- Kroupa P., Bouvier J., 2003, MNRAS, 346, 343
- Kroupa P., Weidner C., Pflamm-Altenburg J., Thies I., Dabringhausen J., Marks M., Maschberger T., 2013, Planets, Stars and Stellar Systems. Volume 5: Galactic Structure and Stellar Populations, Springer Science+Business Media Dordrecht
- Lada C. J., Lada E. A., 2003, ARA&A, 41, 57
- Larsen S. S., 1999, A&AS, 139, 393
- Larsen S. S., 2009, A&A, 494, 539
- Megeath S.T. et al., 2016, AJ, 151, 5
- Meingast S. et al., 2016, A&A, 587, 153
- Parmentier G., Gilmore, G., 2007, MNRAS, 377, 352
- Pflamm-Altenburg J., González-Lópezlira R. A., Kroupa P., 2013, MNRAS, 435, 2604
- Pflamm-Altenburg J., Kroupa P., 2008, Nature, 455, 641
- Porcel C., Garzon F., Jimenez-Vicente J., Battaner E., 1998, A&A, 330, 136
- Randriamanakoto Z., Escala A., Väisänen P., Kankare E., Kotilainen J., Mattila S., Ryder S., 2013, ApJ, 775, L38
- Recchi S., Kroupa P., 2015, MNRAS, 446, 4168
- Schulz C., Pflamm-Altenburg J., Kroupa P., 2015, A&A, 582, 93
- Skibba R. A. et al., 2011, ApJ, 738, 89

Tamburro D., Rix H. W., Walter F., Brinks E., de Blok W. J. G., Kennicutt R. C., Mac Low M. M., 2008, *AJ*, 136, 2872

Weidner C., Kroupa P., Larsen S. S., 2004, *MNRAS*, 350, 1503

Weidner C., Kroupa P., Pflamm-Altenburg J., Vazdekis A., 2013, *MNRAS*, 436, 3309

Yamaguchi R. et al., 2001, *PASJ*, 53, 985

APPENDIX A: THE LECMF WITH A NON-INFINITESIMAL SURFACE AREA

In this section we discuss what differences there are to Sec. 2 if one uses an observable non-infinitesimal surface area ΔA instead of dA . First Eq. (1) becomes

$$\xi_{\text{lecl}}(M_{\text{ecl}}; r) dM_{\text{ecl}} \Delta A = dN_{\text{ecl}}. \quad (\text{A1})$$

For a given region ΔA , $\Sigma_{\text{SFR}}(r)$ becomes the local star formation rate LSFR:

$$\text{LSFR}(r) = \frac{\Delta \text{SFR}}{\Delta A}, \quad (\text{A2})$$

with ΔSFR being the star formation rate in the galactic region ΔA .

The total mass formed in stars is then obtained by multiplying $\text{LSFR}(r)$ with δt (WKL):

$$\text{LSFR}(r) \delta t = \int_{M_{\text{ecl}, \min}}^{M_{\text{U}, \text{loc}}(r)} M'_{\text{ecl}} \xi_{\text{lecl}}(M'_{\text{ecl}}; r) dM'_{\text{ecl}}. \quad (\text{A3})$$

For $\beta \neq 2$ we obtain

$$K(r) = \frac{\text{LSFR}(r) \delta t (2 - \beta)}{M_{\text{U}, \text{loc}}^{2-\beta}(r) - M_{\text{ecl}, \min}^{2-\beta}}. \quad (\text{A4})$$

For the special case of $\beta = 2$,

$$K(r) = \frac{\text{LSFR}(r) \delta t}{\ln(M_{\text{U}, \text{loc}}(r)/M_{\text{ecl}, \min})}. \quad (\text{A5})$$

Using the LECMF it is now possible to calculate the mass of the most massive observable cluster in the region ΔA ($M_{\text{ecl}, \max, \text{loc}}(r)$). To determine $M_{\text{ecl}, \max, \text{loc}}(r)$ we use two conditions: first there is only one most massive cluster. Second the mass of the most massive cluster is $M_{\text{ecl}, \max, \text{loc}}(r)$.

To implement the first condition we choose a mass interval between the upper mass limit $M_{\text{U}, \text{loc}}(r)$ and $M_{\text{ecl}, \text{t}, \text{loc}}(r)$, with $M_{\text{ecl}, \text{t}, \text{loc}}(r)$ chosen in such a way that there is only one cluster between these limits:

$$1 \approx \int_{M_{\text{ecl}, \text{t}}(r)}^{M_{\text{U}, \text{loc}}(r)} \xi_{\text{lecl}}(M'_{\text{ecl}}; r) dM'_{\text{ecl}} \Delta A, \quad (\text{A6})$$

and the second condition implies that the mass between these limits is $M_{\text{ecl}, \max, \text{loc}}(r)$:

$$M_{\text{ecl}, \max, \text{loc}}(r) \approx \int_{M_{\text{ecl}, \text{t}}(r)}^{M_{\text{U}, \text{loc}}(r)} M'_{\text{ecl}} \xi_{\text{lecl}}(M'_{\text{ecl}}; r) dM'_{\text{ecl}} \Delta A. \quad (\text{A7})$$

The reason for the equation being approximated is that $\xi_{\text{lecl}}(M'_{\text{ecl}}; r)$ depends on r . One would have to perform an integration over ΔA to get the exact value (see also Sect. 3). But for small ΔA the above equation is a good approximation.

Using Eqs. (A6) and (A7) $M_{\text{ecl}, \max, \text{loc}}(r)$ becomes for $\beta \neq 2$:

$$M_{\text{ecl}, \text{t}}(r) \approx \left(M_{\text{U}, \text{loc}}^{1-\beta}(r) - \frac{1-\beta}{K(r) \Delta A} \right)^{\frac{1}{1-\beta}}, \quad (\text{A8})$$

$$M_{\text{ecl}, \max, \text{loc}}(r) \approx \frac{K(r)}{2-\beta} [M_{\text{U}, \text{loc}}^{2-\beta}(r) - M_{\text{ecl}, \text{t}}^{2-\beta}(r)] \Delta A. \quad (\text{A9})$$

And for $\beta = 2$:

$$M_{\text{ecl}, \max, \text{loc}}(r) \approx K(r) \left[\ln \left(1 + \frac{M_{\text{U}, \text{loc}}(r)}{K(r) \Delta A} \right) \right] \Delta A. \quad (\text{A10})$$

This is a simple method to determine $M_{\text{ecl}, \max, \text{loc}}(r)$ without having to perform an integration over the area, as long as ΔA is small enough compared to the galaxy.

APPENDIX B: COMPARISON TO THE WKL NORMALIZATION

WKL defined Eq. (4) slightly differently:

$$\Sigma_{\text{SFR}}(r) \delta t = \int_{M_{\text{ecl}, \min}}^{M_{\text{ecl}, \max, \text{loc}}(r)} M'_{\text{ecl}} \xi_{\text{lecl}}(M'_{\text{ecl}}; r) dM'_{\text{ecl}}. \quad (\text{B1})$$

Instead of using a theoretical most massive cluster $M_{\text{U}, \text{loc}}(r)$ as an upper mass limit, $M_{\text{ecl}, \max, \text{loc}}(r)$ was used. This $M_{\text{ecl}, \max, \text{loc}}(r)$ was defined by claiming that there was exactly one cluster in the mass interval $[M_{\text{ecl}, \max, \text{loc}}(r), M_{\text{U}, \text{loc}}(r)]$:

$$1 \approx \int_{M_{\text{ecl}, \max, \text{loc}}(r)}^{M_{\text{U}, \text{loc}}(r)} \xi_{\text{lecl}}(M'_{\text{ecl}}; r) dM'_{\text{ecl}} \Delta A. \quad (\text{B2})$$

A criticism of this LECMF formulation is the claim that Eq. (B2) would result in one most massive cluster with mass $M_{\text{ecl}, \max, \text{loc}}(r)$. In fact it only ensures that there is a most massive cluster without guaranteeing that this most massive cluster has the mass $M_{\text{ecl}, \max, \text{loc}}(r)$. For this another equation is needed:

$$M_{\text{ecl}, \max, \text{loc}}(r) \approx \int_{M_{\text{ecl}, \max, \text{loc}}(r)}^{M_{\text{U}}} M'_{\text{ecl}} \xi_{\text{lecl}}(M'_{\text{ecl}}; r) dM'_{\text{ecl}} \Delta A. \quad (\text{B3})$$

Now it is ensured that there is a mass $M_{\text{ecl}, \max, \text{loc}}(r)$ between $M_{\text{ecl}, \max, \text{loc}}(r)$ and M_{U} . But in general an equation system

$$\begin{aligned} 1 &= \int_a^b f(x) dx, \\ a &= \int_a^b x f(x) dx, \end{aligned} \quad (\text{B4})$$

does not have a real, non-imaginary, solution for power laws. If $f(x)$ is a distribution function this set of equations requests that the mean of x over the interval $[a, b]$ has the same value as the minimal value of x . This can only be possible if $a = b$.

The WKL normalization is thus not correct, but the differences obtained when applying it relative to the correct normalization (Eq. 4) are not significant.

APPENDIX C: OTHER MODELS

Alternatives to the exponential model described in Sec. 3.1:

C1 Phantom Cluster Model

In the past Eq. (B2) had been used to calculate the LECMF. It had been assumed that M_U is a very large mass, often approximated as infinity. In order to get similar results to the previous method we use a slightly modified form of that equation (introduced in Schulz et al. 2015):

$$1 \approx \int_{M_{U,\text{loc}}(r)}^{M_\infty} \xi_{\text{iecl}}(M'_{\text{ecl}}; r) dM'_{\text{ecl}} \Delta A, \quad (\text{C1})$$

with M_∞ going towards infinity.

This resulting model, the phantom cluster model, makes the claim that if there would not be an upper mass limit, then there would be one more cluster in the mass range between the upper mass limit and an infinite mass. A phantom cluster, so to speak. But there is no physical reason for claiming that there is exactly one cluster between the upper mass limit and infinity (Schulz et al. 2015).

The resulting relation between $K(r)$ and $M_{U,\text{loc}}$ is:

$$K(r) \approx (\beta - 1) M_{U,\text{loc}}^{\beta-1}(r) / \Delta A. \quad (\text{C2})$$

If one inserts Eq. (C2) into Eq. (13), so inserting $K(r)$ as a function of $M_{U,\text{loc}}(r)$, a direct relation between $M_{U,\text{loc}}(r)$ and r is obtained:

$$\int_{M_{\text{ecl},\text{min}}}^{M_{U,\text{loc}}(r)} (\beta - 1) \frac{M_{\text{ecl}}^{1-\beta}}{M_{U,\text{loc}}^{\beta-1}(r) \Delta A} dM_{\text{ecl}} \approx \frac{M_{\text{tot}}}{2\pi r_d^2} e^{-r/r_d}. \quad (\text{C3})$$

This equation is not analytically solvable for $M_{U,\text{loc}}$. But numerically it is possible to calculate for every r a corresponding $M_{U,\text{loc}}(r)$ and $K(r)$. The only remaining free parameters are β and ΔA .

We can use these two last equations to calculate a $R'(M_{\text{ecl}})$ for every M_{ecl} , as in the exponential model.

ΔA has to be sufficiently small, so that Eq. (C1) is still a good approximation, but has to be large enough to ensure that a complete LECMF can be found in the area. In the following we make the assumption of ΔA being constant, for simplicity reasons.

The free parameters (β and ΔA) can be fixed by comparing the model to the empirical SFR- $M_{\text{ecl},\text{max}}$ by WKL (see Eq. (23))

Similar to the local case (see Eqs. (A6) and (A7)) the mass of the heaviest cluster in the entire galaxy ($M_{\text{ecl},\text{max}}$) is determined by

$$1 = \int_{M_{\text{ecl},\text{t}}}^{M_U} \xi_{\text{iecl}}(M'_{\text{ecl}}) dM'_{\text{ecl}} \quad (\text{C4})$$

and

$$M_{\text{ecl},\text{max}} = \int_{M_{\text{ecl},\text{t}}}^{M_U} M'_{\text{ecl}} \xi_{\text{iecl}}(M'_{\text{ecl}}) dM'_{\text{ecl}}. \quad (\text{C5})$$

Using the above Eqs. (C4) and (C5) it is possible to calculate for any β a corresponding SFR- $M_{\text{ecl},\text{max}}$ -relation. These relations can be compared with the empirical found relation from Eq. (23) to constraint β . For $\beta = 1.73 \pm 0.01$ and $\Delta A = (2.9 \pm 0.1) \text{ kpc}^2$ the SFR- $M_{\text{ecl},\text{max}}$ -curve aligns best to the fit from WKL. M_U should be larger than any observed cluster. Given the error bars from the observation this requirement is reasonably fulfilled by the chosen parameters, as can be seen in Fig. C1.




Figure C1. The observational data are as in Fig. 2. Plotted here is $M_{\text{ecl},\text{max}}$ and M_U against the SFR as calculated using the phantom cluster model for $\beta = 1.73$ and $\Delta A = 2.9 \text{ kpc}^2$. All observed clusters should be below the M_U -line within their error-bars, which is not the case here but the data might be consistent with this requirement given the uncertainties. See also Schulz et al. (2015) for a discussion of the outlying data points.

Another concern is whether the chosen ΔA is still small enough that Eq. C1 is still a good approximation for an integration over the area ΔA . Numerical tests show that using $\Delta A = 2.9 \text{ kpc}^2$ results in a deviation of up to 3%, compared to an integration over ΔA , which is still quite accurate. Also observations of the LECMF often use a larger observational area, e.g. Lada & Lada (2003) use an area of roughly 12.6 kpc^2 .

C2 Constant K Model

The following is an ansatz that tries to be as simple as possible while being consistent with the observational data. If we assume that K does not depend on the galactocentric distance r , then Eq. (13) should also be valid for $r = 0$. Hence K has for $\beta \neq 2$ the form (with $M_{U,\text{loc}}(0) = M_U$):

$$K = \frac{(2 - \beta) M_{\text{tot}}}{2\pi r_d^2 (M_U^{2-\beta} - M_{\text{ecl},\text{min}}^{2-\beta})} \quad (\text{C6})$$

and for $\beta = 2$:

$$K = \frac{M_{\text{tot}}}{2\pi r_d^2 \ln \left(\frac{M_U}{M_{\text{ecl},\text{min}}} \right)}. \quad (\text{C7})$$

The corresponding $M_{U,\text{loc}}$ as obtained from Eq. (13), using the above form for K , is for $\beta \neq 2$:

$$M_{U,\text{loc}}(r) = \left[e^{-\frac{r}{r_d}} (M_U^{2-\beta} - M_{\text{ecl},\text{min}}^{2-\beta}) + M_{\text{ecl},\text{min}}^{2-\beta} \right]^{\frac{1}{2-\beta}}, \quad (\text{C8})$$

and for $\beta = 2$:

$$M_{U,\text{loc}}(r) = M_{\text{ecl},\text{min}} \left(\frac{M_U}{M_{\text{ecl},\text{min}}} \right)^{e^{-\frac{r}{r_d}}}. \quad (\text{C9})$$

We can use these two last equations to calculate $R'(M_{\text{ecl}})$, as in the exponential model. For $\beta \neq 2$:

$$R'(M_{\text{ecl}}) = -r_d \ln \left[\frac{M_{\text{ecl}}^{2-\beta} - M_{\text{ecl,min}}^{2-\beta}}{M_U^{2-\beta} - M_{\text{ecl,min}}^{2-\beta}} \right] \quad (\text{C10})$$

and for $\beta = 2$:

$$R'(M_{\text{ecl}}) = -r_d \ln \left[\frac{\ln \left(\frac{M_{\text{ecl}}}{M_{\text{ecl,min}}} \right)}{\ln \left(\frac{M_U}{M_{\text{ecl,min}}} \right)} \right]. \quad (\text{C11})$$

Consequently $\xi_{\text{iecl}}(M_{\text{ecl}})$ takes the form

$$\xi_{\text{iecl}}(M_{\text{ecl}}) = K \int_0^{2\pi} \int_0^{R'(M_{\text{ecl}})} M_{\text{ecl}}^{-\beta} r \, dr d\phi. \quad (\text{C12})$$

It is now possible to solve this integral analytically to get an IECMF, which depends only on M_{ecl} :

$$\xi_{\text{iecl}}(M_{\text{ecl}}) = K \pi R'^2(M_{\text{ecl}}) M_{\text{ecl}}^{-\beta}. \quad (\text{C13})$$

This model has two remaining free parameters (β and M_U). As in the other models we want to calculate $M_{\text{ecl,max}}$ in order to compare it to the empirical SFR- $M_{\text{ecl,max}}$ by WKL (see Eq. (23)) and constrain these parameters. Similar to the local case (see Eqs. (A6) and (A7)) the mass of the heaviest cluster in the entire galaxy ($M_{\text{ecl,max}}$) is determined by

$$1 = \int_{M_{\text{ecl,t}}}^{M_U} \xi_{\text{iecl}}(M'_{\text{ecl}}) dM'_{\text{ecl}} \quad (\text{C14})$$

and

$$M_{\text{ecl,max}} = \int_{M_{\text{ecl,t}}}^{M_U} M'_{\text{ecl}} \xi_{\text{iecl}}(M'_{\text{ecl}}) dM'_{\text{ecl}}. \quad (\text{C15})$$

Therefore we can find for every β a SFR- $M_{\text{ecl,max}}$ -curve that aligns with Eq. (23). Doing so we can find for every β a corresponding M_U :

$$M_U = \frac{74138}{2-\beta} \text{SFR}^{0.91-0.15\beta}. \quad (\text{C16})$$

In order to be realistic this M_U has to be heavier than any relevant observed cluster for this specific SFR, but also not so much larger than the mass of the heaviest observed cluster that there would be an unrealistic gap between them. As can be seen in Fig. C2, $\beta = 1.83 \pm 0.1$ fulfils these observational constraints well.

C3 Comparison of the three models

All the models can be written as

$$\xi_{\text{iecl}}(M_{\text{ecl}}) = f(M_{\text{ecl}}, \beta) M_{\text{ecl}}^{-\beta}, \quad (\text{C17})$$

with $f(M_{\text{ecl}}, \beta)$ varying from model to model. All three models use a different $f(M_{\text{ecl}}, \beta)$ and also different values for β in order to be in agreement to the empirical fit by WKL (Eq. 23). Eq. (C17) shows that ξ_{iecl} is not a pure power-law, so β is not the logarithmic slope of ξ_{iecl} . For a comparison of ECMFs resulting from the best fits of the models, see Fig. C3. One can see that the exponential (Sec. 3.1) and the constant K (Sec. C2) model look very similar, although they have different values for β . This is because in these models β is no longer the logarithmic slope. In contrast to the exponential and the constant K model, the phantom cluster model looks different: its logarithmic slope is less steep.




Figure C2. The observational data are as in Fig. 2. Plotted here is $M_{\text{ecl,max}}$ and M_U , as calculated using the constant K model for $\beta = 1.83$ against the SFR. All observed clusters within their uncertainties should be below the M_U -line, which is fulfilled here well.




Figure C3. Comparison of the three models for a galaxy-wide SFR of $1 M_{\odot}/\text{yr}$ and for $\delta t = 10 \text{ Myr}$. For the phantom cluster model $\beta = 1.73$ was used, for the exponential model $\beta = 2.31$ and for the constant K model $\beta = 1.83$. These values have been chosen so that the models fit the observational data shown in Figs. 2, C1 and C2.

The models also result in different dependencies on the distance to the the galactic center in the case of the LECMF. This is illustrated in Fig. C4 with parameters chosen in such a way that the models result in the same LECMF for $r = 0 \text{ kpc}$.

An important difference of the LECMF between the phantom cluster model (Sec. C1) and the other models is that the phantom cluster model has the free parameters β and ΔA , in comparison to the other two which have instead of ΔA a direct dependence on the parameter M_U . Therefore the exponential and the constant K model allow for a M_U which is larger than the masses of the observed clusters (see Figs. 2 and C2).




Figure C4. Comparison of the dependence on the galactocentric distance of the three models. All models are shown with $\beta = 1.73$, $M_U = 400000 M_{\odot}$, $\text{SFR} = 1 M_{\odot}/\text{yr}$, $\delta t = 10 \text{ Myr}$ and $r_d = 2.15 \text{ kpc}$. These values are not the previous fit values but were chosen so that all models produce the same IECMF for a galactocentric distance of 0 kpc. Note that the constant K model leads to a significantly smaller $M_{U,\text{loc}}(r)$, for $r = 5 \text{ kpc}$, than the other models.

Another difference is that the constant K model can be solved analytically, whereas the other models need to be solved numerically.

Summarizing, overall the exponential model, which is physically motivated (Pflamm-Altenburg & Kroupa 2008), works best, as the phantom cluster model depends on a rather arbitrary ΔA and the constant K model does not reproduce well local data. But one may still apply these models, e.g. if one needs an IECMF without numerical modelling, one can use the constant K model.

APPENDIX D: SUMMARY OF VARIABLES

Table D1. List of variables that were introduced.

Variable	Definition	Reason for Introduction
N_{ecl}	number of embedded star clusters	see definition of ξ_{lecl}
M_{ecl}	mass of an embedded star cluster	see definition of ξ_{lecl}
A	surface area	see definition of ξ_{lecl}
r	galactocentric distance	see definition of ξ_{lecl}
ξ_{lecl}	local embedded cluster mass function: $\frac{dN}{dM_{\text{ecl}}dA}$ at r	parameter studied in this paper
K	normalization constant	the normalization constant for ξ_{lecl}
β	power law index	power law index and logarithmic slope of ξ_{lecl}
$M_{\text{ecl,min}}$	minimal embedded cluster mass	the smallest cluster mass
$M_{\text{ecl,max}}$	the largest cluster mass in a given galaxy	used to compare theory with observations
$M_{\text{ecl,max,loc}}$	local $M_{\text{ecl,max}}$	used to compare theory with observations
δt	star formation time-scale	proportional to K
SFR	star formation rate	used to calculate the normalization constant K
LSFR	local star formation rate	proportional to K in the local case
Σ_{SFR}	proportional to K in the local, infinitesimal case	
M_{U}	theoretical most massive cluster physically possible	upper limit for M_{ecl} in a galaxy
$M_{\text{U,loc}}$	local M_{U}	upper limit for M_{ecl} at r
$M_{\text{ecl,t}}$	auxiliary variable	needed together with $M_{\text{U,loc}}$ to calculate $M_{\text{ecl,max,loc}}$: there is exactly one cluster of mass $M_{\text{ecl,max,loc}}$ between $M_{\text{ecl,t}}$ and $M_{\text{U,loc}}$
ξ_{iecl}	integrated embedded cluster mass function : $\frac{dN}{dM_{\text{ecl}}}$	parameter studied in this paper
R_{gal}	radius of the star forming area	upper limit for r
R'	maximal theoretical galactocentric distance for a cluster	necessary to change the r -dependence into a mass dependence
r_d	disk scale length	necessary to describe the exponential galactic disk
$M_{\text{vyc,max}}$	observationally derived maximal very young cluster mass	a good approximation for $M_{\text{ecl,max}}$
WKL	Weidner, Kroupa & Larsen (2004)	gets cited often in this paper